

SMOOTH LYAPUNOV 1-FORMS

M. FARBER, T. KAPPELER, J. LATSCHEV, AND E. ZEHNDER

ABSTRACT. We find conditions which guarantee that a given flow Φ on a closed smooth manifold M admits a smooth Lyapunov one-form ω lying in a prescribed de Rham cohomology class $\xi \in H^1(M; \mathbf{R})$. These conditions are formulated in terms of Schwartzman's asymptotic cycles $\mathcal{A}_\mu(\Phi) \in H_1(M; \mathbf{R})$ of the flow.

1. INTRODUCTION

C. Conley [1, 2] showed that any continuous flow $\Phi : X \times \mathbf{R} \rightarrow X$ on a compact metric space X “*decomposes*” into a chain recurrent flow and a gradient-like flow. More precisely, he proved the existence of a continuous function $L : X \rightarrow \mathbf{R}$ which (i) decreases along any orbit of the flow in the complement $X - R$ of the chain recurrent set $R \subset X$ of Φ and (ii) is constant on the connected components of R . Such a function L is called a *Lyapunov function* for Φ . This existence result plays a fundamental role in Conley's program of understanding general flows as collections of isolated invariant sets linked by heteroclinic orbits.

A more general notion of a *Lyapunov 1-form* was introduced in paper [5]. Lyapunov 1-forms, compared to Lyapunov functions, allow to go one step further and to analyze the flow within the chain-recurrent set R as well. Lyapunov 1-forms provide an important tool in applying methods of homotopy theory to dynamical systems. In the recent papers [4], [5] a generalization of the Lusternik - Schnirelman theory was constructed which applies to flows admitting Lyapunov 1-forms.

The problem of existence of Lyapunov 1-forms was addressed in our recent preprint [6], where we worked in the category of compact metric spaces, continuous flows and continuous closed 1-forms. In the present paper we study the smooth version of the problem: we construct smooth Lyapunov 1-forms for smooth flows on smooth manifolds. We use Schwartzman's asymptotic cycles to formulate a necessary condition for the existence of Lyapunov 1-forms in a given cohomology class. We also show that under an additional

Date: February 27, 2003.

1991 Mathematics Subject Classification. Primary 37B25; Secondary 37C99.

Key words and phrases. Lyapunov function, Lyapunov 1-form, asymptotic cycle.

M. Farber was partially supported by a grant from the Israel Academy of Sciences and Humanities; this work was done while M. Farber visited FIM ETH in Zurich.

T. Kappeler and J. Latschev were partially supported by the European Commission under grant HPRN-CT-1999-00118.

assumption this condition is equivalent to the homological condition introduced in our previous paper [6].

2. DEFINITION

Let V be a smooth vector field on a smooth manifold M . Assume that V generates a continuous flow $\Phi : M \times \mathbf{R} \rightarrow M$ and $Y \subset M$ is a closed, flow-invariant subset.

Definition 1. *A smooth closed 1-form ω on M is called a Lyapunov one-form for the pair (Φ, Y) if it has the following properties:*

- ($\Lambda 1$) *The function $\iota_V(\omega) = \omega(V)$ is negative on $M - Y$;*
- ($\Lambda 2$) *There exists a smooth function $f : U \rightarrow \mathbf{R}$ defined on an open neighborhood U of Y such that*

$$\omega|_U = df \quad \text{and} \quad df|_Y = 0.$$

The above definition is a modification of the notion of a Lyapunov 1-form introduced in section 6 of [5]. The definition of [5] requires that Y consists of finitely many points and the vector field V is locally a gradient of ω with respect to a Riemannian metric.

Definition 1 can also be compared with the definition of a Lyapunov 1-form in the continuous setting which was introduced in [6]. Condition ($\Lambda 1$) above is slightly stronger than condition (L1) of Definition 1 in [6]. Condition ($\Lambda 2$) is similar to condition (L2) of Definition 1 from [6] although they are not equivalent.

There are several natural alternatives for condition ($\Lambda 2$). One of them is:

- ($\Lambda 2'$) *The 1-form ω , viewed as a map $\omega : M \rightarrow T^*(M)$, vanishes on Y .*

It is clear that ($\Lambda 2$) implies ($\Lambda 2'$). We can show that the converse is true under some additional assumptions:

Lemma 1. *If the de Rham cohomology class ξ of ω is integral, $\xi = [\omega] \in H^1(M; \mathbf{Z})$, then the conditions ($\Lambda 2'$) and ($\Lambda 2$) are equivalent.*

Proof. Clearly we only need to show that ($\Lambda 2'$) implies ($\Lambda 2$). Since ξ is integral there exists a smooth map $\phi : M \rightarrow S^1$ such that $\omega = \phi^*(d\theta)$, where $d\theta$ is the standard angular 1-form on the circle S^1 . Let $\alpha \in S^1$ be a regular value of ϕ . Assuming that ($\Lambda 2'$) holds it then follows that $U = M - \phi^{-1}(\alpha)$ is an open neighborhood of Y . Clearly $\omega|_U = df$ where $f : U \rightarrow \mathbf{R}$ is a smooth function which is related to ϕ by $\phi(x) = \exp(if(x))$ for any $x \in U$. Hence ($\Lambda 2$) holds. $\square$

Lemma 2. *The conditions ($\Lambda 2'$) and ($\Lambda 2$) are equivalent if Y is an Euclidean Neighborhood Retract (ENR).*

Proof. Again, we only have to establish ($\Lambda 2'$) $\Rightarrow$ ($\Lambda 2$). Since Y is an ENR it admits an open neighbourhood $U \subset M$ such that the inclusion $i_U : U \rightarrow M$ is homotopic to $i_Y \circ r$, where $i_Y : Y \rightarrow M$ is the inclusion and $r : U \rightarrow Y$ is a retraction (see [3], chapter 4, §8, Corollary 8.7). Pick a base point x_j

in every path-connected component U_j of U and define a smooth function $f_j : U_j \rightarrow \mathbf{R}$ by

$$f_j(x) = \int_{x_j}^x \omega, \quad x \in U_j.$$

The latter integral is independent of the choice of the integration path in U_j connecting x_j with x . This claim is equivalent to the vanishing of the integral $\int_\gamma \omega$ for any closed loop γ lying in U . To show this we apply the retraction to see that γ is homotopic in M to the loop $\gamma_1 = r \circ \gamma$, which lies in Y ; thus we obtain $\int_\gamma \omega = \int_{\gamma_1} \omega = 0$ because of $(\Lambda 2')$. It is clear that the functions f_j together determine a smooth function $f : U \rightarrow \mathbf{R}$ with $df = \omega|_U$. $\square$

Our main goal in this paper is to find topological conditions which guarantee that for a given vector field V on M there exists a Lyapunov 1-form ω lying in a prescribed cohomology class $\xi \in H^1(M; \mathbf{R})$.

3. ASYMPTOTIC CYCLES OF SCHWARTZMAN

Let M be a closed smooth manifold and let V be a smooth vector field. Let $\Phi : M \times \mathbf{R} \rightarrow M$ be the flow generated by V .

Consider a Borel measure μ on M which is invariant under Φ . According to S. Schwartzman [16], these data determine a real homology class

$$\mathcal{A}_\mu = \mathcal{A}_\mu(\Phi) \in H_1(M; \mathbf{R})$$

called *the asymptotic cycle of the flow Φ corresponding to the measure μ* . The class $\mathcal{A}_\mu$ is defined as follows. For a de Rham cohomology class $\xi \in H^1(M; \mathbf{R})$ the evaluation $\langle \xi, \mathcal{A}_\mu \rangle \in \mathbf{R}$ is given by the integral

$$(3.1) \quad \langle \xi, \mathcal{A}_\mu \rangle = \int_M \iota_V(\omega) d\mu,$$

where ω is a closed 1-form in the class ξ . Note that $\langle \xi, \mathcal{A}_\mu \rangle$ is well-defined, i.e. it depends only on the cohomology class ξ of ω , see [16], page 277. Indeed, replacing ω by $\omega' = \omega + df$, where $f : M \rightarrow \mathbf{R}$ is a smooth function, the integral in (3.1) gets changed by the quantity

$$(3.2) \quad \int_M V(f) d\mu = \lim_{s \rightarrow 0} \frac{1}{s} \int_M \{f(x \cdot s) - f(x)\} d\mu(x).$$

Here $V(f)$ denotes the derivative of f in the direction of the vector field V and $x \cdot s$ stands for the flow $\Phi(x, s)$ of the vector field V . Since the measure μ is flow invariant, the integral on the RHS of (3.2) vanishes for any f . It is clear that the RHS of (3.1) is a linear function of $\xi \in H^1(M; \mathbf{R})$. Hence there exists a unique real homology class $\mathcal{A}_\mu \in H_1(M; \mathbf{R})$ which satisfies (3.1) for all $\xi \in H^1(M; \mathbf{R})$.

4. NECESSARY CONDITIONS

We consider the flow Φ as being fixed and we vary the invariant measure μ . As the class $\mathcal{A}_\mu \in H_1(M; \mathbf{R})$ depends linearly on μ , the set of asymptotic cycles $\mathcal{A}_\mu$ corresponding to all Φ -invariant positive measures μ forms a convex cone in the vector space $H_1(M; \mathbf{R})$.

Proposition 1. *Assume that there exists a Lyapunov 1-form for (Φ, Y) lying in a cohomology class $\xi \in H^1(M; \mathbf{R})$. Then*

$$(4.1) \quad \langle \xi, \mathcal{A}_\mu \rangle \leq 0$$

for any Φ -invariant positive Borel measure μ on M ; equality in (4.1) takes place if and only if the complement of Y has measure zero. Further, the restriction of ξ to Y , viewed as a Čech cohomology class

$$\xi|_Y \in \check{H}^1(Y; \mathbf{R})$$

vanishes, $\xi|_Y = 0$.

Proof. Let ω be a Lyapunov one-form for (Φ, Y) lying in the class ξ . According to Definition 1, the function $\iota_V(\omega)$ is negative on $M - Y$ and vanishes on Y . We obtain that the integral

$$\int_M \iota_V(\omega) d\mu = \langle \xi, \mathcal{A}_\mu \rangle$$

is nonpositive.

Assuming $\mu(M - Y) > 0$, we find a compact $K \subset M - Y$ with $\mu(K) > 0$; this follows from the Theorem of Riesz - see e.g. [12], Theorem 2.3(iv), page 256. There is a constant $\epsilon > 0$ such that $\iota_V(\omega)|_K \leq -\epsilon$. Therefore, one has

$$\int_M \iota_V(\omega) d\mu \leq -\epsilon \mu(K) < 0.$$

Hence, the value $\langle \xi, \mathcal{A}_\mu \rangle$ is strictly negative if the measure μ is not supported in Y .

To prove the second statement we observe (see [19]) that the Čech cohomology $\check{H}^1(Y; \mathbf{R})$ equals the direct limit of the singular cohomology

$$\check{H}^1(Y; \mathbf{R}) = \lim_{W \supset Y} H^1(W; \mathbf{R}),$$

where W runs over open neighborhoods of Y . It is clear in view of condition $(\Lambda 2)$ that $\xi|_U = 0 \in H^1(U; \mathbf{R})$ (by the de Rham theorem). Hence the result follows. $\square$

5. CHAIN-RECURRENT SET R_ξ

Given a flow Φ , our aim is to construct a Lyapunov 1-form ω for a pair (Φ, Y) lying in a given cohomology class $\xi \in H^1(M; \mathbf{R})$. A natural candidate for Y is the subset $R_\xi = R_\xi(\Phi)$ of the chain-recurrent set $R = R(\Phi)$ which was defined in [6]. For convenience of the reader we briefly recall the definition.

Fix a Riemannian metric on M and denote by d the corresponding distance function. Given any $\delta > 0$, $T > 1$, a (δ, T) -chain from $x \in M$ to $y \in M$ is a finite sequence $x_0 = x, x_1, \dots, x_N = y$ of points in M and numbers $t_1, \dots, t_N \in \mathbf{R}$ such that $t_i \geq T$ and $d(x_{i-1} \cdot t_i, x_i) < \delta$ for all $1 \leq i \leq N$. Here we use the notation $\Phi(x, t) = x \cdot t$. The chain recurrent set $R = R(\Phi)$ of the flow Φ is defined as the set of all points $x \in M$ such that for any $\delta > 0$ and $T > 1$ there exists a (δ, T) -chain starting and ending at x . The chain recurrent set is closed and invariant under the flow.


Given a cohomology class $\xi \in H^1(M; \mathbf{R})$ there is a natural covering space $p_\xi : \tilde{M}_\xi \rightarrow M$ associated with ξ . A closed loop $\gamma : [0, 1] \rightarrow M$ lifts to a closed loop in $\tilde{M}_\xi$ if and only if the value of the cohomology class ξ on the homology class $[\gamma] \in H_1(M; \mathbf{Z})$ vanishes, $\langle \xi, [\gamma] \rangle = 0$. See [19].

The flow Φ lifts uniquely to a flow $\tilde{\Phi}$ on the covering $\tilde{M}_\xi$. Consider the chain recurrent set $R(\tilde{\Phi}) \subset \tilde{M}_\xi$ of the lifted flow and denote by $R_\xi = p_\xi(R(\tilde{\Phi})) \subset M$ its projection onto M . The set R_ξ is referred to as *the chain recurrent set associated to the cohomology class ξ* . It is clear that R_ξ is a closed and Φ -invariant subset of R . We denote by C_ξ the complement of R_ξ in R ,

$$C_\xi = R - R_\xi.$$

A different definition of R_ξ which does not use the covering space $\tilde{M}_\xi$ can be found in [6].

To state our main result we also need the following notion. A (δ, T) -cycle



of the flow Φ is defined as a pair (x, t) , where $x \in M$ and $t > T$ such that $d(x, x \cdot t) < \delta$. If δ is small enough then any (δ, T) -cycle determines in a canonical way a unique homology class $z \in H_1(M; \mathbf{Z})$ which is represented by the flow trajectory from x to $x \cdot t$ followed by a “short” arc connecting $x \cdot t$ with x . See [6].

6. THEOREM

Theorem 1. *Let V be a smooth vector field on a smooth closed manifold M . Denote by $\Phi : M \times \mathbf{R} \rightarrow M$ the flow generated by V . Let $\xi \in H^1(M; \mathbf{R})$ be a cohomology class such that the restriction $\xi|_{R_\xi}$, viewed as a Čech cohomology*

class $\xi|_{R_\xi} \in \check{H}^1(R_\xi; \mathbf{R})$, vanishes. Then the following properties of ξ are equivalent:

(I). There exists a smooth Lyapunov 1-form for (Φ, R_ξ) in the cohomology class ξ and the subset C_ξ is closed.

(II). For any Riemannian metric on M there exist $\delta > 0$ and $T > 1$ such that the homology class $z \in H_1(M; \mathbf{Z})$ associated with an arbitrary (δ, T) -cycle (x, t) of the flow, with $x \in C_\xi$, satisfies $\langle \xi, z \rangle \leq -1$.

(III). The subset C_ξ is closed and there exists a constant $\eta > 0$ such that for any Φ -invariant positive Borel measure μ on M the asymptotic cycle $\mathcal{A}_\mu = \mathcal{A}_\mu(\Phi) \in H_1(M; \mathbf{R})$ satisfies

$$(6.1) \quad \langle \xi, \mathcal{A}_\mu \rangle \leq -\eta \cdot \mu(C_\xi).$$

(IV). The subset C_ξ is closed and for any Φ -invariant positive Borel measure μ on X with $\mu(C_\xi) > 0$, the asymptotic cycle $\mathcal{A}_\mu = \mathcal{A}_\mu(\Phi) \in H_1(M; \mathbf{R})$ satisfies

$$(6.2) \quad \langle \xi, \mathcal{A}_\mu \rangle < 0.$$

The main point of this result is that it gives sufficient homological conditions for the existence of a Lyapunov 1-form in the cohomology class ξ .

Condition (6.1) can be reformulated using the notion of a quasi-regular point. Recall that $x \in X$ is a *quasi-regular* point of the flow $\Phi : X \times \mathbf{R} \rightarrow X$ if for any continuous function $f : X \rightarrow \mathbf{R}$ the limit

$$(6.3) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t f(x \cdot s) ds$$

exists. It follows from the ergodic theorem that the subset $Q \subset X$ of all quasi-regular points has full measure with respect to any Φ -invariant positive Borel measure on X , see [11], page 106. From the Riesz representation theorem, see e.g. [15], page 256, one deduces that for any quasi-regular point $x \in Q$ there exists a unique positive flow-invariant Borel measure μ_x with $\mu_x(X) = 1$ satisfying

$$(6.4) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t f(x \cdot s) ds = \int_X f d\mu_x$$

for any continuous function f . We use below the well-known fact that any positive, Φ -invariant Borel measure μ with $\mu(X) = 1$ belongs to the weak* closure of the convex hull of the set of measures μ_x , $x \in Q$, see [11], p. 108.

If the subset $C_\xi \subset X$ is closed, and hence compact, one can apply the above mentioned facts to the restriction of the flow to C_ξ . Let ω be an arbitrary smooth closed 1-form lying in the cohomology class ξ . For any quasi-regular point $x \in C_\xi$ of the flow $\Phi|_{C_\xi}$ one has

$$(6.5) \quad \begin{aligned} \lim_{t \rightarrow \infty} \frac{1}{t} \int_x^{x \cdot t} \omega &= \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \iota_V(\omega)(x \cdot s) ds = \\ &= \int_M \iota_V(\omega) d\mu_x = \langle \xi, \mathcal{A}_{\mu_x} \rangle. \end{aligned}$$

We therefore conclude that condition (III) is equivalent to:

(III') *The subset C_ξ is closed and there exists a constant $\eta > 0$ such that for any quasi-regular point $x \in C_\xi$,*

$$(6.6) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \int_x^{x \cdot t} \omega \leq -\eta,$$

where ω is an arbitrary closed 1-form in class ξ .

The value of the limit (6.6) is independent of the choice of a closed 1-form ω ; the only requirement is that ω lies in the cohomology class ξ .

In the special case $\xi = 0$ the set C_ξ is empty and $R = R_\xi$. The above statement then reduces to the following well-known theorem of C.Conley - see [1] and [18], Theorem 3.14:

Proposition 2. (C. Conley) *Let V be a smooth vector field on a smooth closed manifold M . Denote by $\Phi : M \times \mathbf{R} \rightarrow M$ the flow generated by V and by R the chain recurrent set of Φ . Then there exists a smooth Lyapunov function $L : M \rightarrow \mathbf{R}$ for (Φ, R) . This means that $V(L) < 0$ on $M - R$ and $dL = 0$ pointwise on R .*

Proposition 2 is used in the proof of Theorem 1.

As we could not find a proof of this statement in the literature we present one in the appendix.

7. PROOF OF THEOREM 1

The implication (I) $\Rightarrow$ (II) follows from the proof of Proposition 4 in [6].

(II) $\Rightarrow$ (III) By [6], Theorem 2, the set C_ξ is closed. Now we want to show that the inequality (6.2) is satisfied for any positive Φ -invariant Borel measure μ on X with $\mu(C_\xi) > 0$. Fix a closed 1-form ω in the cohomology class ξ . By Lemma 6 from [6], there exist constants $\alpha > 0$ and $\beta > 0$ such that for any $x \in C_\xi$ and $t > 0$, one has

$$\int_x^{x \cdot t} \omega \leq -\alpha t + \beta.$$

Set $t_0 = 2\beta/\alpha$. Then for any $x \in C_\xi$ and $t \geq t_0$ we have

$$(7.1) \quad \frac{1}{t} \int_x^{x \cdot t} \omega \leq -\frac{\alpha}{2}.$$

With any quasi-regular point $x \in C_\xi$ one associates in a canonical way a positive Φ -invariant Borel measure μ_x on C_ξ , see above. It has the property that

$$(7.2) \quad \lim_{t \rightarrow \infty} \frac{1}{t} \int_x^{x \cdot t} \omega = \int_M \iota_V(\omega) d\mu_x.$$

From (7.1) and (7.2) one obtains

$$(7.3) \quad \langle \xi, \mathcal{A}_{\mu_x} \rangle \leq -\frac{\alpha}{2} < 0$$

for any quasi-regular point $x \in C_\xi$. According to [11], page 108, any positive Φ -invariant Borel measure μ with $\mu(M) = \mu(C_\xi) = 1$ belongs to the weak* closure of the convex hull of the set of measures $\{\mu_x; x \in C_\xi \text{ is quasi-regular}\}$; hence

$$(7.4) \quad \langle \xi, \mathcal{A}_\mu \rangle \leq -\frac{\alpha}{2} < 0.$$

It is well known that every (finite) positive Φ -invariant Borel measure is supported on $R = R_\xi \cup C_\xi$, see e.g. [13], Proposition 4.1.18, page 141. As R_ξ and C_ξ are closed and flow-invariant we may write $\mu = \mu_1 + \mu_2$ where μ_1, μ_2 are Φ -invariant and μ_1 is supported on R_ξ , while μ_2 is supported on C_ξ . It follows from (7.4) that $\langle \xi, \mathcal{A}_{\mu_2} \rangle \leq -\frac{\alpha}{2} \cdot \mu_2(C_\xi)$. Further, we claim that $\langle \xi, \mathcal{A}_{\mu_1} \rangle = 0$ for the following reason. Since $\xi|_{R_\xi} = 0$ (as a Čech cohomology class), for any smooth closed 1-form ω on M representing ξ there exists a smooth function f defined on an open neighborhood of R_ξ such that $\omega = df$ near R_ξ . Then we obtain

$$(7.5) \quad \langle \xi, \mathcal{A}_{\mu_1} \rangle = \int_M \iota_V(\omega) d\mu_1 = \int_{R_\xi} \iota_V(\omega) d\mu_1 = \int_{R_\xi} V(f) d\mu_1 = 0.$$

The last equality holds since the measure μ_1 is Φ -invariant (see e.g. [16], Theorem on page 277). Finally, as $\mathcal{A}_\mu = \mathcal{A}_{\mu_1} + \mathcal{A}_{\mu_2}$ we see that $\langle \xi, \mathcal{A}_\mu \rangle \leq -\eta \cdot \mu(C_\xi)$ with $\eta = \alpha/2$ which completes the proof of (II) $\Rightarrow$ (III).

The implication (III) $\Rightarrow$ (VI) is obvious.

We are left to show the implication (IV) $\Rightarrow$ (I). Our argument uses the technique of Schwartzman [16]. It is to show that under the conditions (IV) there exists a smooth Lyapunov 1-form for (Φ, R_ξ) in the class ξ . In a first step we prove that there exists a smooth, closed 1-form ω_1 in the class ξ so that $\iota_V(\omega_1) < 0$ on C_ξ . To this end, denote by $\mathcal{D} \subset C^0(M)$ the space of functions

$$\mathcal{D} = \{V(f); f : M \rightarrow \mathbf{R} \text{ is smooth}\}$$

and by $\mathcal{C}^-$ the convex cone in $C^0(M)$ consisting of all functions $f \in C^0(M)$ with

$$f(x) < 0 \quad \text{for all } x \in C_\xi.$$

As C_ξ is compact, the cone $\mathcal{C}^-$ is open in the Banach space $C^0(M)$ of continuous functions on M , endowed with the usual supremum norm. Choose an arbitrary smooth, closed 1-form ω in the class ξ . Assume that $\mathcal{C}^- \cap (\iota_V(\omega) + \mathcal{D}) = \emptyset$. It then follows from the Hahn - Banach Theorem (cf. [15], page 58) that there exists a continuous linear functional $\Lambda : C^0(M) \rightarrow \mathbf{R}$ so that

$$\Lambda|_{\iota_V(\omega) + \mathcal{D}} \geq 0 \quad \text{and} \quad \Lambda|_{\mathcal{C}^-} < 0.$$

Since $\iota_V(\omega) + \mathcal{D}$ is an affine subspace and Λ is bounded on it from below, we obtain that Λ restricted to $\mathcal{D}$ vanishes. According to the Riesz representation

theorem (cf. [12]), there exists a Borel measure μ on M so that

$$\Lambda(f) = \int_M f d\mu$$

for any $f \in C^0(M)$. By Theorem [16], page 277, the condition $\Lambda|_{\mathcal{D}} = 0$ implies that μ is Φ -invariant. On the other hand, $\Lambda|_{\mathcal{C}^-} < 0$ implies that $\mu|_{C_\xi} > 0$.

Denote by $\chi : M \rightarrow \mathbf{R}$ the characteristic function of C_ξ and let $\nu = \chi \cdot \mu$. As C_ξ is Φ -invariant ν is a Φ -invariant Borel measure and (unlike, possibly, μ) is positive. Note that $\mu - \nu$ is a Φ -invariant Borel measure supported on R_ξ (again using that any Φ -invariant measure is supported on $R = R_\xi \cup C_\xi$). Thus, it follows from our assumption $\xi|_{R_\xi} = 0$, by the same argument which led to (7.5), that

$$\langle \xi, \mathcal{A}_{\mu-\nu} \rangle = 0.$$

Since $\mathcal{A}_{\mu-\nu} = \mathcal{A}_\mu - \mathcal{A}_\nu$ we find

$$\langle \xi, \mathcal{A}_\nu \rangle = \langle \xi, \mathcal{A}_\mu \rangle = \int_M \iota_V(\omega) d\mu = \Lambda(f) \geq 0$$

where $f = \iota_V(\omega)$, contradicting condition (IV). This means that the intersection $\mathcal{C}^- \cap (\iota_V(\omega) + \mathcal{D})$ cannot be empty, i.e. there exists a smooth function $g : M \rightarrow \mathbf{R}$ so that the smooth closed 1-form $\omega_1 = \omega + dg$ is in the class ξ and satisfies

$$\iota_V(\omega_1) < 0 \quad \text{on} \quad C_\xi.$$

This completes the first step of the proof.

To finish the argument, we now adjust ω_1 on the complement of C_ξ so that the resulting form is a Lyapunov 1-form for (Φ, R_ξ) . As $\iota_V(\omega_1) < 0$ on C_ξ and C_ξ is compact, there is some open neighborhood W_1 of C_ξ such that $W_1 \cap R_\xi = \emptyset$ and $\iota_V(\omega_1) < 0$ on W_1 . Since $\xi|_{R_\xi} = 0$, there exists an open neighborhood W_2 of R_ξ such that $W_1 \cap W_2 = \emptyset$ and a smooth function $g : M \rightarrow \mathbf{R}$ such that $\omega_1|_{W_2} = dg$ and $dg|_{W_1} = 0$. By Proposition 2 there exists a smooth Lyapunov function $L : M \rightarrow \mathbf{R}$ for (Φ, R) . Now consider

$$(7.6) \quad \omega_2 = \omega_1 - dg + \lambda dL,$$

where $\lambda > 0$ remains to be chosen. Clearly, the form ω_2 is smooth and closed and represents the class ξ . For any $\lambda > 0$ it satisfies $\omega_2|_{W_2} = d(\lambda L)$, because $\omega_1 - dg$ vanishes on this set. In particular, ω_2 has property (A2) of a Lyapunov 1-form for the pair (Φ, R_ξ) . Note also that for all positive λ we have $\iota_V(\omega_2) < 0$ on W_1 (by the construction of W_1) and on $W_2 - R_\xi$ because $\omega_1 - dg$ vanishes there, whereas $V(L) < 0$. As the complement of $W_1 \cup W_2$ is compact and disjoint from R ,

$$1 < \lambda_0 := 1 + \sup_{x \notin W_1 \cup W_2} \frac{|\iota_V(\omega_1 - dg)|}{|V(L)|} < \infty,$$

and $\iota_V(\omega_2) < 0$ on $M - R_\xi$ for all $\lambda \geq \lambda_0$, showing that for such choices of λ the form ω_2 also has property (A1) of a Lyapunov 1-form for (Φ, R_ξ) .

This completes the proof of the implication (IV) $\Rightarrow$ (I) and hence the proof of Theorem 1. $\square$

APPENDIX: PROOF OF PROPOSITION 2

Recall from [1, II 6.2.A] the alternative characterization of the chain recurrent set R as

$$R = \bigcap \{A \cup A^* \mid (A, A^*) \text{ is an attractor-repeller pair}\}$$

Here a closed, flow-invariant subset $A \subset M$ is called an attractor if it admits a neighborhood U such that A is the maximal flow-invariant subset in the closure of $U \cdot [0, \infty)$. The dual repeller A^* is the set of all points $x \in M$ whose forward limit set is disjoint from A (cf. [1, II 5.1]). Equivalently, (A, A^*) is an attractor-repeller pair if and only if both A and A^* are closed flow invariant subsets of M and the forward (resp. backward) limit set of every point $x \notin A \cup A^*$ is contained in A (resp. A^*) - see [14, Prop. 1.4.].

As M is a closed manifold and hence separable, the number of distinct attractor-repeller pairs is at most countable (cf. [1, II 6.4.A]). Let $\{(A_n, A_n^*)\}_{n \geq 1}$ be some enumeration. For each $n \geq 1$, the construction of Robbin and Salamon (Prop. 1.4. of [14] and the remark following it) yields a smooth function $f_n : M \rightarrow [0, 1]$ with $f_n^{-1}(0) = A_n$, $f_n^{-1}(1) = A_n^*$ and $df_n(V) < 0$ on the complement of $A_n \cup A_n^*$. Let c_n be positive constants such that in a fixed finite atlas of charts all partial derivatives of f_n of order $\leq n$ are bounded pointwise in absolute value by c_n . Then

$$L(x) := \sum_{n=1}^{\infty} \frac{f_n(x)}{2^n c_n}$$

is a smooth function having the required properties. In particular, as for any $n \geq 1$ the differential df_n vanishes on $A_n \cup A_n^*$, the differential of L vanishes on R . $\square$

REFERENCES

- [1] C. Conley, *Isolated invariant sets and the Morse index*, CBMS regional conference series in mathematics, no. 38, AMS, 1976.
- [2] C. Conley, *The gradient structure of a flow: I*, *Ergod. Th. & Dynam. Sys.*, **8**(1988), 11–26
- [3] A. Dold, *Lectures on Algebraic Topology*, Springer - Verlag, 1972.
- [4] M. Farber, *Zeroes of closed 1-forms, homoclinic orbits and Lusternik–Schnirelman theory*, "Topological Methods in Nonlinear Analysis", **19**(2002), 123 - 152
- [5] M. Farber, *Lusternik–Schnirelman Theory and Dynamics*, in: "Lusternik - Schnirelmann Category and related topics", *Contemporary Mathematics*, **316**(2002).
- [6] M. Farber, T. Kappeler, J. Latschev, E. Zehnder, *Lyapunov 1-forms for flows*, to appear in "Ergodic Theory and Dynamical Systems", available also as math.DS/0210473
- [7] J. Franks, *A Variation on the Poincaré-Birkhoff Theorem*, *Contemporary Mathematics*, **81**(1988), 111–117.
- [8] J. Franks, *Homology and dynamical systems*, CBMS regional conference series in mathematics, no. 49, AMS, 1982.
- [9] D. Fried, *The geometry of cross sections to flows*, *Topology*, vol. 21 (1982), 353–371

- [10] F. B. Fuller, *On the surface of section and periodic trajectories*, Amer. J. Math., **87** (1965), 473–480
- [11] K. Jacobs, *Neuere Methoden in der Ergodentheorie*, Springer, 1960
- [12] S. Lang, *Real and Functional Analysis*, third edition, Graduate Texts in Mathematics, 142, 1993, Springer - Verlag
- [13] A. Katok, B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Cambridge University Press, 1995
- [14] J. Robbin and D. Salamon, *Lyapunov maps, simplicial complexes and the Stone functor*, Ergod. Th. & Dynam. Sys. , 12 (1992), pp. 153–183
- [15] W. Rudin, *Functional Analysis*, McGraw-Hill Book Company, 1973
- [16] S. Schwartzman, *Asymptotic cycles*, Ann. of Math., vol. 66, 1957, 270–284
- [17] S. Schwartzman, *Global cross-sections of compact dynamical systems*, Proc. Nat. Acad. Sci. USA **48**(1962), 786 - 791
- [18] M. Shub, *Global Stability of Dynamical Systems*, Springer Verlag, 1986
- [19] E. Spanier, *Algebraic Topology*, Springer Verlag, 1966
- [20] F. W. Wilson, *Smoothing derivatives of functions and applications*, Trans. Amer. Math. **139**(1969), 413 - 428

DEPARTMENT OF MATHEMATICS, TEL AVIV UNIVERSITY, TEL AVIV 69978, ISRAEL
E-mail address: mfarber@tau.ac.il

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ZÜRICH, 8057 ZÜRICH, SWITZERLAND
E-mail address: tk@math.unizh.ch

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ZÜRICH, 8057 ZÜRICH, SWITZERLAND
E-mail address: janko@math.unizh.ch

DEPARTMENT OF MATHEMATICS, ETH ZÜRICH, 8092 ZÜRICH, SWITZERLAND
E-mail address: eduard.zehnder@math.ethz.ch